

# PAN in the Eastern Pacific Free Troposphere: A Satellite View of the Sources, Seasonality, Interannual Variability and Timeline for Trend Detection



Liye Zhu<sup>1</sup>, Emily V. Fischer<sup>1</sup>, Vivienne H. Payne<sup>2</sup>, Thomas Walker<sup>2</sup>, John R. Worden<sup>2</sup>, Zhe Jiang<sup>2,3</sup>, and Susan S. Kulawik<sup>4</sup>  
<sup>1</sup>Department of Atmospheric Science, Colorado State University, Fort Collins, CO, USA. <sup>2</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA. <sup>3</sup>Now at National Center for Atmospheric Research, Boulder, CO, USA. <sup>4</sup>Bay Area Environmental Research Institute Moffett Field, Moffett Field, CA, USA.

## Introduction

- Peroxyacetyl nitrate (**PAN**) is an important reservoir species for NO<sub>x</sub> in the troposphere and plays a significant role in the redistribution of NO<sub>x</sub> to remote regions.
- PAN comprises the dominant fraction of oxidized reactive nitrogen in plumes of Asian origin observed over the Pacific Ocean.
- PAN decomposition in specific plumes of Asian origin subsiding over the **eastern Pacific Ocean** can lead to significant O<sub>3</sub> enhancements.
- Though most past work in this region has focused on **spring**, model calculations by *Jiang et al.* [2016] show that the contribution of PAN to Asian O<sub>3</sub> export is also significant in **summer**.

## TES PAN observations: interannual, seasonal variability


- TES PAN observations**
  - Detection limit ~ 0.2 ppbv.
  - Peak sensitivity 400 – 800 hPa.
  - Only select data with DOFS > 0.6.



Liye Zhu (liyazhu@rams.colostate.edu)  
This work was supported by NASA Award Number NNX14AF14G



## Sources of PAN over eastern Pacific Ocean

- Trans-Pacific transport of PAN clearly also regularly occurs in summer, and potentially at a higher frequency than in spring.



Method:

- 5-day backward HYSPLIT trajectories for each TES PAN retrieval at 4 km;
- Each 1° x 1° grid cell was calculated as the numbers of trajectories passing through a 200 km radius of each latitude and longitude point divided by the total hours of each trajectory and the total number of TES retrievals.



## Timeline for trend detection using TES PAN

- To detect a trend\* of a given magnitude in PAN, it would be **faster** to use satellite observations over the eastern Pacific Ocean region rather than surface in situ observations.
- A trend of a given magnitude will be **more quickly** detected in **summer** than spring in the TES PAN data.

Increasing rate in PAN (% year <sup>-1</sup> )	1	2	3	4	5	6
Years (based on April retrievals)	22	14	11	10	8	8
Years (based on July retrievals)	15	10	8	7	6	6
Years (based on springtime observations at MBO site, [Fischer et al., 2011])	31	20	15	13	11	10

\* A real trend is indicated at the 95% confidence level.

Method:

- 4-day averages of the tropospheric average of all the successfully detected TES PAN retrievals; -> approximately 40 samples for all five years
- Randomly select 8 samples (number for a month) for each year of the trend;
- A hypothetical PAN trend (e.g. a 3% increase in PAN per year) was then added to these selected samples;
- Calculate the correlation coefficient of PAN against time and the t-statistic;
- Repeat this process 1000 times.

